

A new approach to multi-jet calculations in hadron collisions

M. Moretti^a

^aDepartment of Physics, University of Ferrara and INFN sezione di Ferrara

The ALPHA algorithm to evaluate the exact, tree-level matrix elements is reviewed in the context of multi-parton processes in QCD. The algorithm is suited for the automatic calculation of tree-level scattering amplitudes and allows for straightforward inclusion of mass effects. Its CPU cost grows like K^n (K constant, n number of external particles) as opposed to the factorial growth of Feynman graphs. I also discuss the summation over colour configurations which is designed to allow the construction of parton-level event generators suitable to interfacing with a parton-shower evolution including the effects of colour-coherence. Explicit results for the total rates and differential distributions of processes with 8 final-state partons are given.

1. Introduction

The ability to evaluate production rates for multi-jet final states will be fundamental at the LHC to study a large class of processes, within and beyond the SM. A necessary feature of any multi-jet calculation is the possibility to properly evolve the purely partonic final state, for which exact fixed-order perturbative calculations can be performed, into the observable hadronic final state. This evolution is best performed using shower Monte Carlo calculations. The accurate description of color-coherence effects, furthermore, requires a careful bookkeeping of the contribution to the matrix elements of all possible color configurations. The goal of the algorithm [1] described in this talk is to allow the effective calculation of multi-parton matrix elements, allowing the separation, to the leading order in $1/N_c^2$ ($N_c = 3$ being the number of colors), of the independent color configurations. This technique allows an unweighting of the color configurations, and allows the merging of the parton level calculation with the HERWIG Monte Carlo.

The key element of the strategy is the use of the algorithm ALPHA, introduced in Ref. [2] for the evaluation of arbitrary multi-parton matrix elements. The algorithm is built up from an *iteration* of matrix multiplication and is intended for the *automatic* calculation of tree-level amplitudes. It has a complexity growing like a power in the number of particles, compared to

the factorial-like growth that one expects from naive diagram counting.

The ALPHA algorithm will be reviewed in section (2), and its application to multi-jets physics in section (3).

2. The ALPHA algorithm

In reference [2], a new approach to the computation of tree level scattering amplitudes was introduced. This approach, based on the numerical Legendre transform of the effective action, is particularly useful for the automatic calculation of multi-particle processes. This technique was implemented in a FORTRAN code [2] which has been successfully used to study several intricate electroweak processes [3, 4] of interest both at LEP [3] and at the NLC [4]. Among the key features of the algorithm are:

- Suitable (easy to implement) for the automatic calculation of the scattering amplitudes.
- Inclusion of mass effects is straightforward
- The CPU cost grows like K^n (K constant, n number of external particles) as opposed to the factorial growth of the number of Feynman graphs.

We review in this Section the ALPHA algorithm; the interested reader can find a more detailed dis-

cussion in the original paper [1, 2], which includes an explicit analytic example for the $\lambda\phi^3$ theory.

Let Γ be the one-particle-irreducible generator of the Green functions for a given theory. Then the computation of the S-matrix requires the evaluation of the Legendre transform, Z , of Γ :

$$Z(J^\alpha) = -\Gamma(\phi^\alpha) + J^\alpha(x)\phi^\alpha(x)$$

where ϕ^α are the classical fields defined as the solutions of

$$J^\alpha = \frac{\delta\Gamma}{\delta\phi^\alpha},$$

and the J^α play the role of classical sources.

For concreteness we will develop in some detail the case of the scattering amplitude for n external gluons. At tree-level Γ coincide with the Lagrangian. In momentum space

$$\begin{aligned} \mathcal{L}_{YM}(A) = & -1/2(p_\mu A_\nu^a - p_\nu A_\mu^a)^2 \\ & + gf_{abc}(p_\mu A_\nu^a - p_\nu A_\mu^a)A_\mu^b A_\nu^c \\ & - (B_{\mu\nu}^a)^2 - 2gf_{abc}B_{\mu\nu}^a A_\mu^b A_\nu^c + J_\mu^a A_\mu^a \end{aligned}$$

where $B_{\mu\nu}^a$ is an auxiliary field and it is introduced in order to deal with trilinear interactions only. The sources J_μ^a is a standard source term: $J(p) = \sum_{j=1}^n \epsilon_\mu^a \delta(p - p_j)$, *i.e.* it contains the relevant excitation for the external particles. Notice that because of this choice only a finite number of momenta (linear combinations of p_j) enters into the problem which is reduced to a problem with a finite number of degrees of freedom.

The fields $A(p)$ are then found as solutions of the equation of motion. Let us stick to Feynman gauge for definiteness.

$$\begin{aligned} A_\mu^a(p) = & \frac{g}{p^2} f_{abc} [2(k-p) \cdot A^b(q) A_\mu^c(k) \\ & - 2q_\mu A^b(q) \cdot A_\mu^c(k) \\ & - B_{\mu\nu}^b(q) A_\nu^c(k)] + \frac{1}{p^2} J_\mu(p) \\ B_{\mu\nu}^a(p) = & -gf_{abc} A_\mu^b(q) A_\nu^c(k) \end{aligned} \quad (1)$$

where momentum conservation is understood.

The equations of motion (1) are solved *iteratively* (expansion in g) and this implies that the

problem is solved with a *loop of matrices multiplication*, more suitable for numerical implementation than the standard approach. The initialization steps are

$$\begin{aligned} A_\mu^a(p_j) &= \epsilon_\mu^a(p_j) \\ B_{\mu\nu}^a(p_j) &= 0 \end{aligned} \quad (2)$$

and the subsequent steps

$$\begin{aligned} A_\mu^a(p_j + p_k) = & \frac{gf_{abc}}{(p_j + p_k)^2} [-2(p_j + 2p_k) \cdot A^b(p_j) A_\mu^c(p_k) \\ & + 2(p_k + 2p_j) \cdot A^b(p_k) A_\mu^c(p_j) \\ & - 2(p_j + p_k)_\mu A^b(p_k) \cdot A^c(p_j) \\ & - B_{\mu\nu}^b(p_j) A_\mu^c(p_k)] \\ B_{\mu\nu}^a(p_j + p_k) = & -gf_{abc} A^b(p_k) A_\mu^c(p_j) \end{aligned} \quad (3)$$

The step (3) is then iterated until we have constructed A and B fields with up to n momenta. An important remark is in order here. Performing the iteration (3) we drop terms containing $\epsilon^a(p_j)\epsilon^a(p_j)$ *i.e.* twice the same external momenta. In fact, although legitimate, they don't contribute to the amplitude.

Finally the amplitude is

$$\mathcal{A}(p_{j_1}, \dots, p_{j_n}) = \mathcal{L}_{YM}(A) \quad (4)$$

where the A fields are obtained in eqns. (2,3)¹. Notice that the prescription to drop terms containing at least twice the same external momenta ($\sim \epsilon^a(p_j)\epsilon^a(p_j)$) is still kept and it is because of this prescription together with the choice (2) for the initial step that neither truncation nor functional derivation is actually required.

A remark is in order here. The algorithm sketched above has an important feature: it provides a very compact way of storing the relevant information. Indeed the number of contraction $A^a(p_{j_1} + \dots + p_{j_m})$ (see eq. 3) which is needed is of the order 2^n , each of them requiring roughly the

¹Using the equation of motion it is possible to halve the required number of iteration step as well as to reduce the number of contributions entering the final expression of the amplitude.

same CPU time to be computed. Therefore the CPU cost of the algorithm grows like a *constant to the power n* , n being the number of external particles, as opposed to the factorial growth of the number of Feynman graphs. To hint how does this work let us develop in more detail the case of the amplitude for five external gluons. For simplicity we neglect the four gluons coupling. The A fields are given by

$$\begin{aligned}
A_\mu^a(p_j) &= \epsilon_\mu^a(p_j) \quad j = 1, \dots, 5 \\
A_\mu^a(p_j + p_k) &= \\
&\frac{gf_{abc}}{(p_j + p_k)^2} \left[-2(p_j + 2p_k) \cdot A^b(p_j) A_\mu^c(p_k) \right. \\
&\quad \left. + 2(p_k + 2p_j) \cdot A^b(p_k) A_\mu^c(p_j) \right. \\
&\quad \left. - 2(p_j + p_k)_\mu A^b(p_k) \cdot A^c(p_j) \right] \\
&\quad j = 1, \dots, 5
\end{aligned} \tag{5}$$

and the ALPHA amplitude by

$$\begin{aligned}
\mathcal{A} &= g \sum_{j=1}^5 A^a(p_j)_\alpha \\
&\sum_{\substack{k \neq l \neq m \neq n \neq j}}^5 f_{abc} \tau_{\alpha\beta\gamma} A^b(p_k + p_l)_\beta A^c(p_m + p_n)_\gamma \\
\tau_{\alpha\beta\gamma} &= g_{\alpha\beta}(p_j - p_k - p_l)_\gamma \\
&\quad + g_{\gamma\alpha}(p_m + p_n - p_j)_\beta \\
&\quad + g_{\beta\gamma}(p_k + p_l - p_m - p_n)_\alpha
\end{aligned} \tag{6}$$

notice that the summations over k, l, m, n , as well as over lorentz and color indices, are carried over *before* the multiplication by $A^a(p_j)$. It is this feature extensively used in the construction of both the A fields and the amplitude which turns the factorial growth of the number of Feynman graphs to a power law growth of the CPU cost of the ALPHA algorithm. Notice that in eqns. (5,6) $A(p_j + p_k + p_m)$ and $A(p_j + p_k + p_m + p_l)$ are not computed (and used). Moreover neither the kinetic term nor the source one appears in the expression (6) of the amplitude. These simplifications occur because of the equation of motion as discussed in more detail in [1, 2].

3. Multi-jets processes

Multi-jet final states play an important role in the study of high-energy collisions. They provide in fact interesting signatures for several phenomena, both within the Standard Model (e.g. top-pair production), and beyond it (e.g. multi-jet decays of supersymmetric particles such as gluinos and squarks). The accurate determination of the properties of these phenomena requires a good understanding of the properties of the usually large multi-jet QCD backgrounds, which can distort the shapes of signal distributions and affect the measurement of quantities such as resonances' masses.

There are several reasons for wanting to improve the tools currently available to perform these calculations.

1. First of all, interesting final states with larger jet multiplicities will become available with the next generation of colliders (LHC and NLC).
2. Secondly, one would like to be able to complement the calculation of parton-level matrix elements with the evaluation of the full hadronic structure of the final state.

The goal of the algorithm [1] described in this Section is to allow the effective calculation of multi-parton matrix elements, allowing the separation, to the leading order in $1/N_c^2$ ($N_c = 3$ being the number of colors), of the independent color configurations. This technique allows an unweighting of the color configurations, and allows the merging of the parton level calculation with the HERWIG Monte Carlo. The key ingredient of our strategy is the ALPHA algorithm outlined in the previous section. It's power-law growing in complexity is a necessary feature of any attempt to evaluate matrix elements for processes with large numbers of external particles, since the number of Feynman diagrams grows very quickly (see table 1) beyond any reasonable value.

Once the hard scattering matrix element is known the prescription [1] to correctly generate the parton-shower associated to a given event in the large- N_c limit is the following:

Process	$n = 7$	$n = 8$	$n = 9$	$n = 10$
$g g \rightarrow n g$	559,405	10,525,900	224,449,225	5,348,843,500
$q\bar{q} \rightarrow n g$	231,280	4,016,775	79,603,720	1,773,172,275

Table 1

Number of Feynman diagrams corresponding to amplitudes with different numbers of quarks and gluons.

1. Calculate the $(n - 1)!$ dual amplitudes corresponding to all possible planar color configurations. This can be done with **ALPHA** using N_c large enough [1].
2. Extract the *most likely* color configuration for this event on a statistical basis, according to the relative contribution of the single configurations to the total event weight ². Since each dual amplitude is gauge invariant, the choice of color-configurations is also a gauge-invariant operation.
3. Develop the PS out of each initial and final-state parton, starting from the selected color configuration. This step can be carried out by feeding the generated event to a Monte-Carlo program such as **HERWIG**, which is precisely designed to *turn partons into jets* starting from an assigned color-ordered configuration.

Notice that, if the dual amplitudes are evaluated for a specific helicity configuration, **HERWIG** will also include spin-correlation effects in the evolution of the parton shower [5, 6, 7, 8, 9].

As a result, use of the dual-amplitude representation of a multi-gluon amplitude allows to accurately describe not only the large-angle correlations in multi-jet final states, but also the full shower evolution of the initial- and final-state partons with the same accuracy available in **HERWIG** for the description of 2-jet final states.

In alternative to the above prescription, one can use **ALPHA** to calculate the matrix elements

²Defining $w_i = |A_i|^2$ for each color flow i , and $W_i = \sum_{k=1, \dots, i} w_k / \sum_{k=1, \dots, n} w_k$, the j -th color structure will be selected if $W_{j-1} \leq \xi < W_j$, for a random number ξ uniformly distributed over the interval $[0, 1]$.

for external states with assigned colors. Since these states are all orthogonal, such an approach is particularly efficient if one wants to use a Monte Carlo method to sum over all possible color states. The program will then extract through a standard unweighting (at the leading order in $1/N_c^2$) a specific color flow from all possible color flows contributing to a given orthogonal color state. This color flow is then suitable as an initial condition for the shower evolution. The advantage of this approach is twofold: first the number of dual amplitudes contributing to the amplitude for external states with assigned colors is substantially smaller than the total one and second dual amplitudes are required only for *accepted events* which, in general, are a small fraction of the generated ones. Further details can be found in [1].

3.1. Results

As an example of our technique, we present here results for the following two parton-level processes:

$$\begin{aligned}
 g g &\rightarrow 8 g \\
 q \bar{q} &\rightarrow 8 g.
 \end{aligned}
 \tag{7}$$

For comparisons, we also computed the above reactions in the simple approximation first suggested by Kunszt and Stirling [10]. This approximation (hereafter referred to as **SPHEL**) consists in assuming that the average value of maximally helicity violating amplitudes [11] is equal to the average value of all other non-zero amplitudes.

The kinematic configuration and the cut values used in our numerical examples are as follows:

$$\begin{aligned}
 \sqrt{s} &= 1500 \text{ GeV}, & p_{T_i} &> 60 \text{ GeV}, & |\eta_i| &< 2, \\
 \Delta R_{ij} &> 0.7.
 \end{aligned}$$

These values, and the choice of a fixed strong cou-




Figure 1. Differential distributions for the minimum gluon transverse momentum. Exact result vs. SPHEL.

pling $\alpha_s = 0.12$, only serve for illustrative purposes.

In Fig. 1, we show the distribution of the minimum gluon transverse momentum for both processes. In Fig 2 we plot the distributions for the maximum gluon transverse momentum.

4. Conclusions

We have reviewed the **ALPHA** algorithm to evaluate the exact, tree-level matrix elements in the context of multi-parton processes in QCD. The algorithm is suitable (easy to implement) for the automatic calculation of the scattering amplitudes and it allows the inclusion of mass effects in a straightforward manner. The CPU cost of the algorithm grows like K^n (K constant, n number of external particles) as opposed to the factorial growth of the number of Feynman graphs.

This technique has been tested for processes such as $gg \rightarrow n$ gluons and $q\bar{q} \rightarrow n$ gluons, with n up to 9. We discussed how the summation over




Figure 2. Differential distributions for the maximum gluon transverse momentum. Exact result vs. SPHEL.

colour configurations allows the construction of parton-level event generators suitable to interfacing with a parton-shower evolution including the effects of colour-coherence. This will eventually lead to a fully exclusive, hadron-level description of multi-jet final states, accurately incorporating the dynamics of large jet-jet separation angles.

REFERENCES

1. F. Caravaglios, M. L. Mangano, M. Moretti and R. Pittau, Nucl. Phys. **B539** (1999) 215
2. F. Caravaglios and M. Moretti, Phys. Lett. **B358** (1995) 332
3. F. Caravaglios and M. Moretti, Z. Phys. **C74** (1997) 291;
G. Montagna, M. Moretti, O. Nicrosini and F. Piccinini, Nucl. Phys. **B541** (1999) 31;
G. Montagna, M. Moretti, O. Nicrosini, A. Pallavicini and F. Piccinini, Nucl. Phys. **B547** (1999) 39

4. M. Moretti, Nucl. Phys. **B484** (1997) 3;
G. Montagna M. Moretti, O. Nicrosini and
F. Piccinini, *Eur.Phys.J.* **C2** (1998) 483;
F. Gangemi, G. Montagna, M. Moretti,
O. Nicrosini and F. Piccinini, Eur. Phys. J.
C9 (1999) 31;
F. Gangemi, G. Montagna, M. Moretti,
O. Nicrosini and F. Piccinini, Nucl. Phys.
B559 (1999) 3
5. J. C. Collins, Nucl. Phys. **B304** (1988) 794.
6. I. G. Knowles, Nucl. Phys. **B310** (1988) 571.
7. G. Marchesini and B. R. Webber, Nucl. Phys.
B310 (1988) 461.
8. G. Marchesini, B. R. Webber, G. Abbiendi,
I. G. Knowles, M. H. Seymour and L. Stanco,
Comput. Phys. Commun. **67** (1992) 465.
9. G. Marchesini, B. R. Webber, G. Abbiendi,
I. G. Knowles, M. H. Seymour and L. Stanco,
hep-ph/9607393.
10. Z. Kunszt and W.J. Stirling, Phys. Rev. **D37**
(1988) 2439.
11. S.J. Parke and T. Taylor, Phys. Rev. Lett. **56**
(1986) 2459.